


# Determination of $\alpha_s$ from the QCD static energy

Antonio Vairo

Technische Universität München



## Bibliography

- (1) A. Bazakov, N. Brambilla, X. Garcia i Tormo, P. Petreczky, J. Soto and A. Vairo  
*Determination of  $\alpha_s$  from the QCD static energy: an update*  
Phys. Rev. D90 (2014) 7, 074038 arXiv:1407.8437
- (2) X. Garcia i Tormo  
*Review on the determination of  $\alpha_s$  from the QCD static energy*  
Mod. Phys. Lett. A28 (2013) 1330028 arXiv:1307.2238
- (3) A. Bazakov, N. Brambilla, X. Garcia i Tormo, P. Petreczky, J. Soto and A. Vairo  
*Determination of  $\alpha_s$  from the QCD static energy*  
Phys. Rev. D86 (2012) 114031 arXiv:1205.6155
- (4) N. Brambilla, X. Garcia i Tormo, J. Soto and A. Vairo  
*Precision determination of  $r_0 \Lambda_{\overline{\text{MS}}}$  from the QCD static energy*  
Phys. Rev. Lett. 105 (2010) 212001 arXiv:1006.2066
- (5) N. Brambilla, X. Garcia i Tormo, J. Soto and A. Vairo  
*The QCD static energy at NNNLL*  
Phys. Rev. D80 (2009) 034016 arXiv:0906.1390
- (6) N. Brambilla, X. Garcia i Tormo, J. Soto and A. Vairo  
*The logarithmic contribution to the QCD static energy at  $N^4\text{LO}$*   
Phys. Lett. B647 (2007) 185 arXiv:hep-ph/0610143


$\alpha_s$  in 2014

# Status of $\alpha_s$

	$\alpha_s(M_Z^2)$	
Alekhin [2001]	$0.1143 \pm 0.013$	DIS [4]
BBG [2004]	$0.1134^{+0.0019}_{-0.0021}$	valence analysis, NNLO [5,6]
GRS	0.112	valence analysis, NNLO [7]
ABKM	$0.1135 \pm 0.0014$	HQ: FFNS $N_f = 3$ [8]
JR14	$0.1136 \pm 0.0004$	dynamical approach [9]
JR14	$0.1162 \pm 0.0006$	including NLO-jets [9]
MSTW	$0.1171 \pm 0.0014$	(2009) [10]
Thorne	0.1136	[DIS+DY, HT*] (2014) [11]
ABM11 <sub>J</sub>	$0.1134 - 0.1149 \pm 0.0012$	Tevatron jets (NLO) incl. [12]
ABM13	$0.1133 \pm 0.0011$	[13]
ABM13	$0.1132 \pm 0.0011$	(without jets) [13]
CTEQ	$0.1159 \dots 0.1162$	[14]
CTEQ	0.1140	(without jets) [14]
NN21	$0.1174 \pm 0.0006 \pm 0.0001$	[15]
Gehrmann et al.	$0.1131^{+0.0028}_{-0.0022}$	$e^+e^-$ thrust [16]
Abbate et al.	$0.1140 \pm 0.0015$	$e^+e^-$ thrust [17]
CMS	$0.1151 \pm 0.0033$	$t\bar{t}$ [18]
NLO Jets ATLAS	$0.111^{+0.0017}_{-0.0007}$	[19]
NLO Jets CMS	$0.1148 \pm 0.0055$	[19]
BBG [2004]	$0.1141^{+0.0020}_{-0.0022}$	valence analysis, N <sup>3</sup> LO [5,6]
3-jet rate	$0.1175 \pm 0.0025$	Dissertori et al. 2009 [20]
Z-decay rate	$0.1189 \pm 0.0026$	BCK 2008/12 (N <sup>3</sup> LO) [21,22]
$\tau$ -decay rate	$0.1212 \pm 0.0019$	BCK 2008 (N <sup>3</sup> LO) [21,22]
$\tau$ -decay rate	$0.1204 \pm 0.0016$	Pich 2011 [1]
$\tau$ -decay rate	$0.325 \pm 0.018$ (at $m_\tau$ )	FOTP: [23]
$\tau$ -decay rate	$0.374 \pm 0.025$ (at $m_\tau$ )	CIPT: [23]
Lattice	$0.1205 \pm 0.0010$	PACS-CS 2009 (2+1 fl.) [24]
Lattice	$0.1184 \pm 0.0006$	HPQCD 2010 [25]
Lattice	$0.1200 \pm 0.0014$	ETMC 2012 (2+1+1 fl.) [26]
Lattice	$0.1156 \pm 0.0022$	Bazavov et al. (2+1 fl.) [27]
Lattice	$0.1130 \pm 0.0010(stat)$	RBC-UKQCD (preliminary, 2014) [28]

## PDG average

Current uncertainties of  $\alpha_s$  are not fully reflected in the PDG average.



# Theory

## Static energy

$$E_0(r) = \lim_{T \rightarrow \infty} \frac{i}{T} \ln \langle \boxed{\quad} \rangle; \quad \boxed{\quad} = \exp \left\{ ig \oint dz^\mu A_\mu \right\}$$

Perturbation theory describes  $E_0(r)$  in the **short range** ( $r\Lambda \ll 1$ ,  $\alpha_s(1/r) < 1$ ):

$$E_0(r) = \Lambda_s - \frac{C_F \alpha_s}{r} (1 + \# \alpha_s + \# \alpha_s^2 + \# \alpha_s^3 + \# \alpha_s^3 \ln \alpha_s + \# \alpha_s^4 \ln^2 \alpha_s + \# \alpha_s^4 \ln \alpha_s + \dots)$$

- $E_0(r)$  is known at **three loops**.
  - Anzai Kiyo Sumino PRL 104 (2010) 112003  
A.Smirnov V.Smirnov Steinhauser PRL 104 (2010) 112002
- $\ln \alpha_s$  signals the cancellation of contributions coming from **different energy scales**:


$$\ln \alpha_s = \ln \frac{\mu}{1/r} + \ln \frac{\alpha_s/r}{\mu}$$

- Brambilla Pineda Soto Vairo PRD 60 (1999) 091502

## Energy scales


In the short range the static Wilson loop is characterized by a hierarchy of energy scales:

$$1/r \gg V_o - V_s \gg \Lambda; \quad V_s \approx -C_F \frac{\alpha_s}{r}, \quad V_o \approx \frac{1}{2N} \frac{\alpha_s}{r}$$



# Effective Field Theories

It is convenient to factorize the contributions from the different scales with EFTs:



$$E_0(r) = \Lambda_s + V_s(r, \mu) - i \frac{g^2}{N} V_A^2 \int_0^\infty dt e^{-it(V_o - V_s)} \langle \text{Tr } \mathbf{r} \cdot \mathbf{E}(t) \mathbf{r} \cdot \mathbf{E}(0) \rangle(\mu) + \dots$$

res. mass	potential	ultrasoft contribution
-----------	-----------	------------------------

- Brambilla Pineda Soto Vairo NPB 566 (2000) 275

The  $\mu$  dependence cancels between

$$V_s \sim \ln r\mu, \ln^2 r\mu, \dots$$

**ultrasoft contribution**  $\sim \ln(V_o - V_s)/\mu, \ln^2(V_o - V_s)/\mu, \dots \ln r\mu, \ln^2 r\mu, \dots$

## Static singlet potential and energy at N<sup>3</sup>LL

$$V_s(r, \mu) = V_s(r, 1/r) - \frac{C_F C_A^3}{6\beta_0} \frac{\alpha_s^3(1/r)}{r} \left\{ \left( 1 + \frac{3}{4} \frac{\alpha_s(1/r)}{\pi} a_1 \right) \ln \frac{\alpha_s(1/r)}{\alpha_s(\mu)} \right. \\ \left. \left( \frac{\beta_1}{4\beta_0} - 6c \right) \left[ \frac{\alpha_s(\mu)}{\pi} - \frac{\alpha_s(1/r)}{\pi} \right] \right\}$$

Summed to the ultrasoft contribution at two loops, it provides the static energy at N<sup>3</sup>LL.

## Mass renormalon

The perturbative expansion of  $V_s$  is affected by a renormalon ambiguity of order  $\Lambda$ . This ambiguity does not affect the slope of the potential (and the extraction of  $\alpha_s$ ).

It may be eliminated from the perturbative series

- either by subtracting a (constant) series in  $\alpha_s$  to  $V_s$  and reabsorb it in a redefinition of the residual mass,
- or by considering the **force**:

$$F(r, \alpha_s(\nu)) = \frac{d}{dr} E_0(r, \alpha_s(\nu))$$

- The force  $F(r, \alpha_s(1/r))$  could be directly compared with lattice,
- or integrated and compared with the static energy

$$E_0(r) = \int_{r_*}^r dr' F(r', \alpha_s(1/r'))$$

up to an irrelevant constant fixed by the overall normalization of the lattice data.

Note that there are no  $\ln \nu r$  ( $\nu$  = renormalization scale).

# Analysis

## Lattice

We use 2+1-flavor lattice QCD obtained from tree-level improved gauge action and Highly-Improved Staggered Quark (HISQ) action by the HotQCD collaboration.  $m_s$  was fixed to its physical value, while  $m_l = m_s/20$ .

This corresponds to a pion mass of about 160 MeV in the continuum limit.

$\beta$	7.373	7.596	7.825
$r_1/a$	5.172(34)	6.336(56)	7.690(58)
Volume	$48^3 \times 64$	$64^4$	$64^4$

The largest gauge coupling,  $\beta = 7.825$ , corresponds to lattice spacings of  $a = 0.041$  fm.

- Bazakov et al PRD 90 (2014) 094503

The lattice spacing was fixed using the  $r_1$  scale defined as  $r^2 \frac{dE_0(r)}{dr} \Big|_{r=r_1} = 1.0$ ;

$r_1 = 0.3106 \pm 0.0017$  fm from the pion decay constant  $f_\pi$ .


- Bazakov et al PoS LATTICE 2010 (2010) 074

## Procedure


We use data for each value of the lattice spacing separately, and at the end perform an average of the different obtained values of  $\alpha_s$  with the following procedure.

- Perform fits to the lattice data for the static energy  $E_0(r)$  at different orders of perturbative accuracy. The parameter of the fits is  $\Lambda_{\overline{\text{MS}}}$ .
- Repeat the above fits for each of the following distance ranges:  $r < 0.75r_1$ ,  $r < 0.7r_1$ ,  $r < 0.65r_1$ ,  $r < 0.6r_1$ ,  $r < 0.55r_1$ ,  $r < 0.5r_1$ , and  $r < 0.45r_1$ .
- Use ranges where the reduced  $\chi^2$  either decreases or does not increase by more than one unit when increasing the perturbative order, or is smaller than 1.
- To estimate the **perturbative uncertainty** of the result, repeat the fits
  - by varying the scale in the perturbative expansion, from  $\nu = 1/r$  to  $\nu = \sqrt{2}/r$  and  $\nu = 1/(\sqrt{2}r)$ ,
  - by adding/subtracting a term  $\pm(C_F/r^2)\alpha_s^{n+2}$  to the expression at  $n$  loops.Take the largest uncertainty.

## Data ranges




$$\chi^2/\text{d.o.f.} \text{ for } \beta = 7.825$$




Fits for  $r < 0.6r_1$  are acceptable. In the final result we will use only fits for  $r < 0.5r_1$ . The fitting curve has been normalized on the 7th, 8th and 9th lattice point respectively.

## $a\Lambda_{\overline{\text{MS}}}$ at different orders of perturbative accuracy for $\beta = 7.825$




# $r_1 \Lambda_{\overline{\text{MS}}}$ at three-loop accuracy




The band shows the determination of 2012.

# Statistical error vs perturbative error




The **statistical error** is estimated by taking values of  $\Lambda_{\overline{\text{MS}}}$  at one  $\chi^2$  unit above minimum.

# Short-distance points vs long-distance points



The band shows the determination of 2012.

## Looking for condensates



By repeating the fits adding a monomial term proportional to  $r^3$  and  $r^2$ , which could be associated with gluon and quark local condensates, and also a term proportional to  $r$ , we do not find evidence for a significant non-perturbative term at short distances and the value of  $\Lambda_{\overline{\text{MS}}}$  remains unchanged.

# Results



## $\Lambda_{\overline{\text{MS}}}$

Results at three-loop plus leading-ultrasoft resummation for the  $r < 0.5r_1$  fit range.  
The final result is the weighted average of different  $\beta$ s with linearly added errors.

	$a\Lambda_{\overline{\text{MS}}}; N_{\text{ref}} = 7$	$a\Lambda_{\overline{\text{MS}}}; N_{\text{ref}} = 8$	$a\Lambda_{\overline{\text{MS}}}; N_{\text{ref}} = 9$	$a\Lambda_{\overline{\text{MS}}}; \text{range spanned}$	$r_1\Lambda_{\overline{\text{MS}}}; \text{range spanned}$
$\beta = 7.373$	$0.0957^{+0.0046}_{-0.0028} \pm 0.0017$	$0.0957^{+0.0046}_{-0.0028} \pm 0.0017$	$0.0957^{+0.0046}_{-0.0028} \pm 0.0017$	$0.0957^{+0.0046}_{-0.0028} \pm 0.0017$	$0.4949^{+0.0240}_{-0.0144} \pm 0.0086 \pm 0.0025$ $= 0.4949^{+0.0256}_{-0.0170}$
$\beta = 7.596$	$0.0781^{+0.0046}_{-0.0029} \pm 0.0007$	$0.0784^{+0.0043}_{-0.0027} \pm 0.0010$	$0.0785^{+0.0046}_{-0.0029} \pm 0.0007$	$0.0783^{+0.0048}_{-0.0031} \pm 0.0010$	$0.4961^{+0.0303+0.0066}_{-0.0197-0.0061} \pm 0.0044$ $= 0.4961^{+0.0313}_{-0.0211}$
$\beta = 7.825$	$0.0644^{+0.0032}_{-0.0019} \pm 0.0006$	$0.0642^{+0.0033}_{-0.0020} \pm 0.0008$	$0.0643^{+0.0032}_{-0.0020} \pm 0.0008$	$0.0643^{+0.0033}_{-0.0021} \pm 0.0008$	$0.4944^{+0.0256}_{-0.0159} \pm 0.0065 \pm 0.0037$ $= 0.4944^{+0.0267}_{-0.0175}$
<b>Average</b>					$r_1\Lambda_{\overline{\text{MS}}} = 0.495^{+0.028}_{-0.018}$



$$r_1\Lambda_{\overline{\text{MS}}} = 0.495^{+0.028}_{-0.018} \quad \text{which converts to} \quad \Lambda_{\overline{\text{MS}}} = 315^{+18}_{-12} \text{ MeV}$$

## Static energy vs lattice data



Note the agreement between perturbation theory and lattice data up to about 0.2 fm.

# Static energy at different perturbative orders vs lattice data



Lattice data with  $\beta$  from 6.664 to 7.825 are displayed.

The red error bars correspond to the errors of the lattice data (include normalization).

$\alpha_s$


$$\alpha_s(1.5 \text{ GeV}, n_f = 3) = 0.336^{+0.012}_{-0.008}$$

which corresponds to

$$\alpha_s(M_Z, n_f = 5) = 0.1166^{+0.0012}_{-0.0008}$$

from four-loop running,  $m_c = 1.6 \text{ GeV}$  and  $m_b = 4.7 \text{ GeV}$ .

## Comparison with other determinations



For  $\tau$  decays (ALEPH + OPAL) see also

$$\alpha_s(M_Z, n_f = 5) = 0.1165 \pm 0.0012 \text{ (FOPT)}, \quad \alpha_s(M_Z, n_f = 5) = 0.1185 \pm 0.0015 \text{ (CIPT)}$$